

Mechanism for spectral break in cosmic ray proton spectrum from Supernova remnant W44

M.A. Malkov¹, P.H. Diamond¹ and R.Z. Sagdeev²

¹CASS and Department of Physics, University of California, San Diego, La Jolla, CA 92093 and

²University of Maryland, College Park, Maryland 20742-3280, USA

Recent observations of the supernova remnant W44 by the *Fermi* spacecraft observatory strongly support the idea that the bulk of galactic cosmic rays is accelerated in such remnants by a Fermi mechanism, also known as diffusive shock acceleration. However, the W44 expands into weakly ionized dense gas, and so a significant revision of the mechanism is required. In this paper we provide the necessary modifications and demonstrate that strong ion-neutral collisions in the remnant surrounding lead to the steepening of the energy spectrum of accelerated particles by *exactly one power*. The spectral break is caused by Alfvén wave evanescence leading to the fractional particle losses. The gamma-ray spectrum generated in collisions of the accelerated protons with the ambient gas is also calculated and successfully fitted to the Fermi Observatory data. The parent proton spectrum is best represented by a classical test particle power law $\propto E^{-2}$, steepening to E^{-3} at $E_{br} \approx 7\text{GeV}$ due to deteriorated particle confinement.

Ongoing *direct* observations of a number of galactic supernova remnants (SNRs) in the X- and gamma-ray bands supported by the radio, optical and UV data [1, 2] seem to rapidly close in on the century long problem of the cosmic ray (CR) origin. Overall, the observations favor the diffusive shock acceleration (DSA, operating in SNR shocks) as a mechanism for the production of galactic CRs. However, there are at least two critical questions that observations pose to the theory. First, what exactly is the form of the spectrum that the theory predicts? Second, when do we really see the proton (i.e., the primary CR component) emission and when do we see just a contaminating but radiatively more efficient electron component?

Several decades of studies of the background galactic CRs, preceding direct observations (that became available only recently) demonstrated that already a simple test particle version of the DSA (leading to a power-law $\propto E^{-2}$ particle energy distribution) reproduces the form of the energy spectrum reasonably well. It is possible, however, that a somewhat steeper power-law would better accommodate the $E^{-2.7}$ measured spectrum [3, 4] (see also the Discussion section below), even though this difference can be attributed to CR propagation losses (also not completely certain). Nevertheless, the *direct observations* of emission coming from particles accelerated in SNR shocks often indicate significantly steeper than E^{-2} spectra [5]. Also the recent *Fermi* Large Area Telescope (LAT) observations of the SNR W44 suggest that the spectrum of the gamma ray producing protons is substantially steeper at high energies than the DSA predicts [6]. It should be noted that a similar discrepancy has already been found in the high energy gamma ray spectrum of the SNR RXJ 1713 measured by the CANGAROO [7] and HESS [5] atmospheric Cerenkov telescopes. The simple DSA prediction was even used as an argument against the hadronic origin of the gamma emission on the ground that if the high energy part of the spectrum was extrapolated (with the DSA index 2) to lower energies, the emission there would be unacceptably high, e.g. [8]. Motivated by this inconsistency with the DSA theory in particular, we have previously published a suggestion of the mechanism for the spectral break [9], which can resolve the discrepancies. We believe that a similar phenomenon is also applicable to the recent observations of the SNR W44.

In particular, we show below that ion-neutral collisions in a molecular cloud adjacent to the remnant steepen the energy spectrum of accelerated particles. We calculate the gamma-ray spectrum resulting from the collisions of accelerated protons with the molecular cloud material and fit the result to the Fermi Observatory data.

Results

Mechanism for the spectral break The physics of the spectral break is very simple. When a SNR shock approaches a molecular cloud (MC) or a pre-supernova swept-up shell, confinement of accelerated particles generally deteriorates. Due to the particle interaction with magnetic turbulence, confinement requires scales similar to the particle gyroradius [10, 11]. While the waves are in a strongly ionized (closer to the shock) medium they propagate freely in a broad frequency range at the Alfvén speed $V_A = B/\sqrt{4\pi\rho_i}$ with the frequencies $\omega = kV_A$. Here k is the wave number (assumed parallel to the local field \mathbf{B}) and ρ_i is the ion mass density. As long as the Alfvén wave frequency is higher than the ion-neutral collision frequency ν_{in} , the waves are weakly damped. When, on the other hand, the ion-neutral collision frequency is higher (deeper into the cloud), neutrals are entrained by the oscillating plasma and the Alfvén waves are also able to propagate, albeit with a factor $\sqrt{\rho_i/\rho_0} < 1$ lower speed, where ρ_0 is the neutral density. The propagation speed reduction occurs because every ion is now “loaded” with ρ_0/ρ_i neutrals. In between these two regimes Alfvén waves are heavily damped and even disappear altogether for sufficiently small $\rho_i/\rho_0 \ll 0.1$. The evanescence wave number range is then bounded by $k_1 = \nu_{in}/2V_A$ and $k_2 = 2\sqrt{\rho_i/\rho_0}\nu_{in}/V_A$. These phenomena have been studied in detail in [12, 13], and specifically in the context of the DSA in [14–16]. Now we turn to their impact on the particle confinement and emissivity.

In the frame work of a quasilinear wave-particle interaction the wave number k is approximately related to the parallel (to the magnetic field) component of the particle momentum p_{\parallel} by the cyclotron resonance condition $kp_{\parallel}/m = \pm\omega_c$ where the (non-relativistic) gyro-frequency $\omega_c = eB/mc$. Therefore, the

frequency range where the waves cannot propagate may be conveniently translated into the parallel momentum range

$$p_1 < |p_{\parallel}| < p_2 \quad (1)$$

with

$$p_1 = 2V_A m \omega_c / v_{in}, \quad p_2 = \frac{p_1}{4} \sqrt{\rho_0 / \rho_i} > p_1, \quad (2)$$

That a spectral break must form at the photon energy corresponding to the particle momentum $p = p_1 = p_{br}$, can be readily understood from Fig.1. The 'dead zones' $p_1 < |p_{\parallel}| < p_2$ imply that particles with $|p_{\parallel}| > p_1$ do not turn around (while moving along the magnetic field) and escape from the region of CR-dense gas collisions at a p_{\parallel}/p fraction of the speed of light. This happens in the region of enhanced gamma radiation. Therefore, an initially isotropic distribution of accelerated particles fills only a slab in momentum space $|p_{\parallel}| < p_1$ and becomes highly anisotropic (a 'pancake' distribution). What matters for the integral emission, however, is a locally isotropic component \bar{f} of this new proton distribution. It can be introduced by re-averaging the 'pancake' ($|p_{\parallel}| < p_1$) distribution in pitch angle, $\bar{f}(p) \equiv \int_0^1 f(p, \mu) d\mu$, and is readily obtained assuming that particles remaining in the dense gas (those with $|p_{\parallel}| < p_1$) maintain their isotropic pitch-angle distribution, i.e.

$$\bar{f}(p) = \int_0^{\mu_1} f_0(p) d\mu = \begin{cases} (p_1/p) f_0(p), & p \geq p_1 \\ f_0(p), & p < p_1 \end{cases} \quad (3)$$

where $f_0(p)$ is the initial (isotropic) distribution function and μ is the cosine of the pitch angle (see Fig.1), $\mu_1 = \min\{p_1/p, 1\}$. Thus, the slope of the particle momentum distribution becomes steeper by exactly one power above $p = p_1 \equiv p_{br}$. In particular, any power-law distribution $\propto p^{-q}$, upon entering an MC, turns into p^{-q-1} at $p \geq p_{br}$, and preserves its form at $p < p_{br}$.

Now the question is whether particle escape from the MC can quench the acceleration process itself. In principle it can, since also the particles from the interval $|p_{\parallel}| < p_1$ cannot return to the shock and continue acceleration. Instead, by slowly leaking to the part of the momentum space with $|p_{\parallel}| > p_1$, they should ultimately escape as well. This would certainly be the case if the MC were filling the entire shock precursor. However, MCs are known to be clumpy [17–19], and may fill only a small fraction of the precursor. In this case the acceleration process continues almost unimpeded, the accelerated protons illuminate the 'cloudlets' and make them visible in γ -rays due to the high density target material.

Break momentum While the one power spectral break in the pitch-angle averaged particle distribution seems to be a robust environmental signature of a weakly ionized medium into which the accelerated particles propagate, the break momentum remains uncertain. According to eq.(2), $p_{br} (\equiv p_1)$ depends on the magnetic field strength and ion density as well

as on the frequency of ion-neutral collisions, $v_{in} = n_0 \langle \sigma V \rangle$. Here $\langle \sigma V \rangle$ is the product of the collision cross-section and collision velocity averaged over the thermal distribution. Using an approximation of [15, 20] for $\langle \sigma V \rangle$, p_{br} can be estimated as

$$p_{br}/mc \simeq 10 B_{\mu}^2 T_4^{-0.4} n_0^{-1} n_i^{-1/2}. \quad (4)$$

Here the gas temperature T_4 is measured in the units of $10^4 K$, magnetic field B_{μ} -in microgauss, n_0 and n_i (number densities corresponding to the neutral/ion mass densities ρ_0 and ρ_i) -in cm^{-3} . Note that the numerical coefficient in the last expression may vary depending on the average ion and neutral masses and can be higher by a factor of a few for typical molecular cloud conditions [12, 21] than the estimate in eq.(4) suggests. The remaining quantities in the last formula are also known too poorly to make an accurate independent prediction of the position of the break in the gamma ray emission region. Those are the regions near the blast wave where complicated physical processes unfold. They include particle acceleration, strong MHD turbulence (driven by particles and their interaction with ambient gas inhomogeneities), gas ionization by shock generated UV photons, turbulent plasma heating and even evaporation of magnetic cloudlets [20, 22, 23]. Also important may be the ionization by the low energy CRs accelerated at the blast wave. However, as their diffusion length is shorter than that of the particles with $p \gtrsim p_{br}$, we may assume that they do not reach the MC. Pre-ionization by the UV photons can also be ignored for the column density $N > 10^{19} cm^{-2}$ ahead of the shock beyond which they are absorbed [24]. The authors or the Ref. [24], using the earlier data from [25] have also analyzed the parameters involved in eq.(4) and found the above estimate of p_{br} to be in a good agreement with the spectral break position measured by the *Fermi* LAT. Nevertheless, we may run the argument in reverse and use the *Fermi* observations [6] of the gamma-ray spectrum of SNR W44 to determine the break momentum in the parent particle spectrum and constrain the parameters in eq.(4). Once we also know the amount of the slope variation Δq , we can calculate the full spectrum up to the cut-off energy.

Particle spectra To calculate the particle spectra, we need to determine the degree of nonlinear modification of the shock structure. In principle, it can be calculated consistently, given the shock parameters and the particle maximum momentum, p_{max} . In the case of a broken spectrum, p_{br} plays the role of p_{max} , since this is the momentum where the dominant contribution to the pressure of accelerated particles comes from, thus setting the scale of the modified shock precursor. Note that in the conventional nonlinear (NL) acceleration theory, the cut-off momentum p_{max} plays this role, because the nonlinear spectra are sufficiently flat so as to make the pressure diverge with momentum.

The break in the photon spectrum is observed at about 2 GeV, which places the break on the proton distribution at about $p_{br} \simeq 7 GeV/c$ [6]. For the strength of the break $\Delta q = 1$, the spectrum above it is clearly pressure converging, and we perform the calculation of the shock structure and the spectrum using this break momentum as the point of the maximum in the CR partial pressure. Once this point is set, we can use

an analytic approach for a stationary nonlinear acceleration problem using p_{br} as an input parameter.

Apart from p_{br} , the nonlinear solution depends on a number of other parameters, such as the injection rate of thermal particles into acceleration, Mach number, the precursor heating rate and the shock velocity V_s . Of these parameters only the latter is known accurately, $V_s \approx 300 \text{ km/s}$, the other parameters are still difficult to ascertain. Fortunately, in sufficiently strong shocks the solution generally tends to either stay close to the test particle (TP) solution (leaving the shock structure only weakly modified) or else it transitions to a strongly modified NL-solution regime. The TP regime typically establishes in cases of moderate Mach numbers, low injection rates and low p_{max} (now p_{br}), while the NL regime is unavoidable in the opposite part of the parameter space.

In the TP regime the spectrum is close to a power-law with the spectral index 2 throughout the supra-thermal energy range. In the NL regime, however, the spectrum develops a concave form, starting from a softer spectrum at the injection energy, with the index $q \simeq (r_s + 2)/(r_s - 1) > 2$, where $r_s < 4$ is the sub-shock compression ratio. Then it hardens, primarily in the region $p \sim mc$, where both the partial pressure and diffusivity of protons change their momentum scaling. The slope reaches its minimum at the cut-off (break) energy, which, depending on the degree of nonlinearity, can be as low as 1.5 or even somewhat lower if the cut-off is abrupt. The question now is into which of these two categories the W44 spectrum falls? Generally, in cases of low maximum (or, equivalently, low spectral break $p_{br} \lesssim 10$) momentum, the shock modification is weak, so the spectrum is more likely to be in the only slightly nonlinear, almost TP regime. On the other hand, there is a putative indication from the electron radio emission that their spectrum may be close to $q_e \approx 1.75$, which could be the signature of a moderately nonlinear acceleration process. It should be remembered, however, that this is a global index across the W44 remnant. There are resolved bright filaments where a canonical $\alpha = -0.5$ spectrum, corresponding precisely to the TP parent electron spectrum with $q_e = 2$ is observed [26]. Moreover, there are regions with the positive indices $\alpha \lesssim 0.4$ which cannot be indicative of a DSA process without corrections for subsequent spectral transformations such as an absorption by thermal electrons [26]. These regions can very well contribute to the overall spectral hardening discussed above, mimicking the acceleration nonlinearity. Finally, secondary electrons give rise to the flattening of the radio spectrum as well [24].

If the accelerated protons and electrons respond to the turbulence similarly, which is almost certainly the case in the ultra-relativistic regime, their spectra should have similar slopes there (as long as the synchrotron losses are ignorable). In using the electron radio spectrum as a probe for the level of acceleration nonlinearity, the following two relations are useful. First, there is a relation between the electron energy and the radio frequency $\nu_{MHz} = 4.6 \cdot B_\mu E_{GeV}^2$. The second relation, $q_e = 1 - 2\alpha$, links the spectral index of radio emission α (assuming the radio flux $\propto \nu^\alpha$) and the spectral index of the parent electrons q_e (assuming their energy spectrum $\propto E^{-q_e}$). Once the global radio spectral index of W44, $\alpha \simeq -0.37$ [26]

is generated by freshly accelerated electrons in the frequency range $74 < \nu < 10700$ MHz, the electrons should maintain their modified spectrum over the energy range spanning more than one order of magnitude. For example, assuming $B_\mu \simeq 70$ [6], one sees that electrons must maintain an index $q_e \approx 1.75$ between $0.46 < E < 5.8$ GeV. While the upper bound is acceptable given the spectral break proton energy inferred from the super GeV emission measured by the Fermi LAT, the lower end is rather uncomfortable, since the nonlinear modification of both protons and electrons with the Bohm (or other similar for protons and electrons turbulent diffusivities) starts (slowly) only at the proton rest energy. The calculated nonlinear spectra are shown in Fig.2 for the both species. At and below 1 GeV, the electron spectrum is very close to the test particle solution, $q_e \approx 2$, even though the proton spectrum may be somewhat steeper there, as we mentioned above.

Photon spectra The above considerations somewhat weaken the radio data as a probe for the slope of the electron and (more importantly) for the proton spectrum. Therefore, the exact degree of nonlinearity of the acceleration remains unknown and we consider both the TP and weakly NL regimes in our calculations of the photon spectra, generated in $p-p$ collisions. The results are shown in Fig.3. The best fit to the Fermi data is provided by a TP energy distribution ($\propto E^{-2}$) below $p_{br} \simeq 7 \text{ GeV/c}$ with the spectrum steepening by exactly one power above it. Note that the small deviation of the computed spectrum from the lowest energy *Fermi*-point may also be corrected by including the Bremsstrahlung of the secondary electrons [24]. The spectrum steepening is perfectly consistent with the proton partial escape described above (with no parameters involved) and shown in Fig.1. For comparison, a weakly NL spectrum, shown in Fig.2, is also used for these calculations (dashed line in Fig.3), but its fit would require a stronger break ($\Delta q > 1$) or a low momentum cut-off, Fig.3, i.e. at least one additional free parameter.

Discussion

To summarize the results, the previously suggested mechanism for a break on the spectrum of shock accelerated protons [9] is found consistent with the recent [6] Fermi LAT observations of the SNR W44. The observed gamma ray spectrum most likely results from the decay of π^0 -mesons which are born in $p-p$ collisions of shock accelerated protons with an ambient dense gas. The parent proton spectrum is best represented by a classical test particle power law $\propto E^{-2}$, steepening to E^{-3} at $E_{br} \approx 7 \text{ GeV}$ due to deteriorated particle confinement caused by the ion-neutral collisions and the resultant Alfvén wave evanescence. The position of the break momentum in the particle spectrum may be estimated using eq.(4), or conversely, the combination of parameters involved in this estimate can be inferred from the measured break momentum. The cut-off momentum is not constrained in this scenario.

An alternative explanation, based on a different mechanism of the break, associated with the change of the particle transport in the CR shock precursor [27] is also possible but is less definitive in the spectrum slope variation Δq across the break

(see also [24] for the most recent alternative suggestions). In addition, the ref [27] mechanism would imply a considerable nonlinearity, i.e. a stronger CR shock precursor than it seems appropriate for the observed low break momentum. Still alternatively, assuming our “environmental” break mechanism is at work, i.e. $\Delta q = 1$, but the shock structure appreciably modified, we arrive at the $E^{-1.75}$ spectrum below the break (as the radio observations suggest for the electrons), and $E^{-2.75}$ above the break. A fit to the data is marginally possible, but it would require a relatively low cut-off momentum at about 100 GeV/c. This possibility may be supported or ruled out once the data (upper limit) around this energy become available.

The most robust and attractive aspect of the suggested mechanism for the spectral break is the exact $\Delta q = 1$ variation of the spectral index. Indeed, this change in the spectral slope is due to truncation of the particle momentum phase space and does not depend on any parameters. In a combination with the test particle regime operating below the break, which is consistent with the low value of the break momentum, it provides a very good fit to the *Fermi* LAT data. Of three physically different types of spectral breaks suggested earlier [9, 27, 28], namely the current, “environmental” mechanism appears to be plausible in situations where dense target gas is present, as required for efficient π^0 production. In a more general context, a spectral break is a natural resolution of the well known but puzzling trend of the *nonlinear* (i.e. supposedly improved) DSA theory to develop spectra which are even harder than the simple test particle spectra, thus encountering more difficulties while accommodating the bulk of observations [3, 4]. Such a spectrum – i.e., diverging in energy – exhausts the shock energy available for the acceleration as the cut-off momentum grows. Note, that the spectrum of the RX J1713.7-3946 [5] is also consistent with the same break mechanism but with $p_{br} \sim 10^3 \text{ GeV}/c$ and with naturally stronger acceleration nonlinearity [9]. However, it is difficult to make the case for hadronic origin of the gamma-ray emission of the RX J1713.7-3946 [5, 29, 30]. The fundamental role of the W44 for the problem of CR origin is that this particular remnant seems to rule out contaminating electron emission due to Bremsstrahlung and the inverse Compton scattering [6, 24] thus favoring the hadronic origin of the gamma emission.

Methods

Acceleration model and proton spectrum Methods of calculation of the accelerated particle spectra, including the par-

ticle back-reaction on the shock structure, are now well developed. We use the diffusion-convection equation for describing the distribution of high energy particles (CRs). To include the back-reaction, three further relations are used in a quasi-stationary acceleration regime. These are the conservation of mass and momentum fluxes in the smooth part of the shock transition (so called CR-precursor) and the Rankine-Hugoniot relation for the shock compression at the gaseous discontinuity (sub-shock). The complete system of equations is then reduced to one nonlinear integral equation [31]. This equation is solved numerically and the results are shown in Fig.2. The above method of calculation of the parent particle spectra is shown to be in excellent agreement [32] with numerical simulations [33, 34] as well as with other semi-analytic approaches [35].

Gamma-ray emission spectrum Once the parent proton spectrum is obtained, we calculate the π^0 production rate and the gamma-ray emissivity. In so doing, we adopt numerical recipe described in detail in [36, 37]. The physical processes behind these calculations are [38] (i) collisions of accelerated protons with the protons of the ambient gas resulting in the following spectrum of π^0 -mesons:

$$F_{pp}(E_\pi) = 4\pi N_{pg} \int \frac{d\sigma(E_\pi, E_p)}{dE_\pi} J_p(E_p) dE_p \quad (5)$$

where, N_{pg} is the number density of protons in the gas, $d\sigma/dE_\pi$ is the differential cross section for the π^0 production in collisions between accelerated protons of energy E_p and gas protons, J_p is the flux of accelerated protons, E_π is the energy of π^0 mesons; (ii) decay of π^0 resulting in the gamma emission spectrum

$$F(E_\gamma) = 2 \int_{E_\gamma + m_\pi^2 c^4 / 4E_\gamma}^{\infty} \frac{F_{pp}(E_\pi)}{\sqrt{E_\pi^2 - m_\pi^2 c^4}} dE_\pi \quad (6)$$

where m_π is the pion rest mass.

-
- [1] Reynolds, S. P. Supernova Remnants at High Energy. *ARA&A* **46**, 89–126 (2008).
 - [2] Raymond, J. C. Optical and UV Diagnostics of Supernova Remnant Shocks. *Space Science Reviews* **99**, 209–218 (2001).
 - [3] Gaisser, T. K., Protheroe, R. J. & Stanev, T. Gamma-Ray Production in Supernova Remnants. *Astrophys. J.* **492**, 219–227 (1998).
 - [4] Hillas, A. M. TOPICAL REVIEW: Can diffusive shock accel-

eration in supernova remnants account for high-energy galactic cosmic rays? *Journal of Physics G Nuclear Physics* **31**, 95–131 (2005).

- [5] Aharonian, F. *et al.* A detailed spectral and morphological study of the gamma-ray supernova remnant RX J1713.7-3946 with HESS. *Astronomy and Astrophysics*. **449**, 223–242 (2006).
- [6] Abdo, A. A. *et al.* Gamma-Ray Emission from the Shell of Supernova Remnant W44 Revealed by the Fermi LAT. *Science*

- 327, 1103–1106 (2010).
- [7] Enomoto, R. *et al.* The acceleration of cosmic-ray protons in the supernova remnant RX J1713.7-3946. *Nature* **416**, 823–826 (2002).
- [8] Reimer, O. & Pohl, M. No evidence yet for hadronic TeV gamma-ray emission from SNR RX J1713.7-3946. *Astronomy and Astrophys.* **390**, L43–L46 (2002).
- [9] Malkov, M. A., Diamond, P. H. & Sagdeev, R. Z. On the Gamma-Ray Spectra Radiated by Protons Accelerated in Supernova Remnant Shocks near Molecular Clouds: The case of Supernova Remnant RX J1713.7-3946. *Astrophys. J. Lett.* **624**, L37–L40 (2005).
- [10] Drury, L. O. An introduction to the theory of diffusive shock acceleration of energetic particles in tenuous plasmas. *Reports on Progress in Physics* **46**, 973–1027 (1983).
- [11] Blandford, R. & Eichler, D. Particle Acceleration at Astrophysical Shocks - a Theory of Cosmic-Ray Origin. *Phys. Rep.* **154**, 1–75 (1987).
- [12] Kulsrud, R. & Pearce, W. P. The Effect of Wave-Particle Interactions on the Propagation of Cosmic Rays. *Astrophys. J.* **156**, 445–469 (1969).
- [13] Zweibel, E. G. & Shull, J. M. Confinement of cosmic rays in molecular clouds. *Astrophys. J.* **259**, 859–868 (1982).
- [14] Voelk, H. J., Morfill, G. E. & Forman, M. A. The effect of losses on acceleration of energetic particles by diffusive scattering through shock waves. *Astrophys. J.* **249**, 161–175 (1981).
- [15] Drury, L. O. C., Duffy, P. & Kirk, J. G. Limits on diffusive shock acceleration in dense and incompletely ionised media. *Astronomy and Astrophys.* **309**, 1002–1010 (1996).
- [16] Reville, B., Kirk, J. G., Duffy, P. & O’Sullivan, S. Environmental Limits on the Nonresonant Cosmic-Ray Current-Driven Instability. *International Journal of Modern Physics D* **17**, 1795–1801 (2008).
- [17] Crutcher, R. M. Magnetic Fields in Molecular Clouds: Observations Confront Theory. *Astrophys. J.* **520**, 706–713 (1999).
- [18] Chevalier, R. A. & Oishi, J. Cassiopeia A and Its Clumpy Pre-supernova Wind. *Astrophys. J. Lett.* **593**, L23–L26 (2003).
- [19] Parizot, E., Marcowith, A., van der Swaluw, E., Bykov, A. M. & Tatischeff, V. Superbubbles and energetic particles in the Galaxy. I. Collective effects of particle acceleration. *Astronomy and Astrophys.* **424**, 747–760 (2004).
- [20] Draine, B. T. & McKee, C. F. Theory of interstellar shocks. *ARA&A* **31**, 373–432 (1993).
- [21] Nakano, T. Contraction of magnetic interstellar clouds. *Fundamentals of Cosmic Physics* **9**, 139–231 (1984).
- [22] Shull, J. M. & McKee, C. F. Theoretical models of interstellar shocks. I - Radiative transfer and UV precursors. *Astrophys. J.* **227**, 131–149 (1979).
- [23] Bykov, A. M., Chevalier, R. A., Ellison, D. C. & Uvarov, Y. A. Nonthermal Emission from a Supernova Remnant in a Molecular Cloud. *Astrophys. J.* **538**, 203–216 (2000).
- [24] Uchiyama, Y., Blandford, R. D., Funk, S., Tajima, H. & Tanaka, T. Gamma-ray Emission from Crushed Clouds in Supernova Remnants. *Astrophys. J. Lett.* **723**, L122–L126 (2010).
- [25] Reach, W. T., Rho, J. & Jarrett, T. H. Shocked Molecular Gas in the Supernova Remnants W28 and W44: Near-Infrared and Millimeter-Wave Observations. *Astrophys. J.* **618**, 297–320 (2005).
- [26] Castelletti, G., Dubner, G., Brogan, C. & Kassim, N. E. The low-frequency radio emission and spectrum of the extended SNR W44: new VLA observations at 74 and 324 MHz. *Astronomy and Astrophys.* **471**, 537–549 (2007).
- [27] Malkov, M. A. & Diamond, P. H. Nonlinear Shock Acceleration Beyond the Bohm Limit. *Astrophys. J.* **642**, 244–259 (2006).
- [28] Malkov, M. A., Diamond, P. H. & Jones, T. W. On the Possible Reason for Nondetection of TeV Protons in Supernova Remnants. *Astrophys. J.* **571**, 856–865 (2002).
- [29] Aharonian, F. A. *et al.* High-energy particle acceleration in the shell of a supernova remnant. *Nature* **432**, 75–77 (2004).
- [30] Katz, B. & Waxman, E. In which shell-type SNRs should we look for gamma-rays and neutrinos from P P collisions? *Journal of Cosmology and Astro-Particle Physics* **1**, 1–29 (2008).
- [31] Malkov, M. A. & Drury, L. O. Nonlinear theory of diffusive acceleration of particles by shock waves. *Reports on Progress in Physics* **64**, 429–481 (2001).
- [32] Moskalenko, I. V., Porter, T. A., Malkov, M. A. & Diamond, P. H. Hadronic Gamma Rays from Supernova Remnants. In *International Cosmic Ray Conference*, vol. 2 of *International Cosmic Ray Conference*, 763–766 (2008).
- [33] Berezhko, E. G. & Ellison, D. C. A Simple Model of Nonlinear Diffusive Shock Acceleration. *Astrophys. J.* **526**, 385–399 (1999).
- [34] Ellison, D. C., Berezhko, E. G. & Baring, M. G. Nonlinear Shock Acceleration and Photon Emission in Supernova Remnants. *Astrophys. J.* **540**, 292–307 (2000).
- [35] Blasi, P., Gabici, S. & Vannoni, G. On the role of injection in kinetic approaches to non-linear particle acceleration at non-relativistic shock waves. *Mon. Not. R. Astron. Soc.* **361**, 907–918 (2005).
- [36] Kamae, T., Karlsson, N., Mizuno, T., Abe, T. & Koi, T. Parameterization of gamma, e^{+/-}, and Neutrino Spectra Produced by p-p Interaction in Astronomical Environments. *Astrophys. J.* **647**, 692–708 (2006).
- [37] Karlsson, N. & Kamae, T. Parameterization of the Angular Distribution of Gamma Rays Produced by p-p Interaction in Astronomical Environments. *Astrophys. J.* **674**, 278–285 (2008).
- [38] Dermer, C. D. Secondary production of neutral pi-mesons and the diffuse galactic gamma radiation. *Astronomy and Astrophys.* **157**, 223–229 (1986).

Acknowledgments

We would like to thank Sabrina Casanova for furnishing a detailed information about the code developed in ref.[37]. Support by NASA under the Grants NNX 07AG83G and NNX09AT94G as well as by the Department of Energy, Grant No. DE-FG02-04ER54738 is gratefully acknowledged.

Author contributions

M.A.M. carried out the calculations of the spectra, P.H.D. contributed to the wave propagation part, R.Z.S. contributed to the wave-particle interaction part. The article was written by M.A.M and edited by all the co-authors.

Corresponding author

M.A. Malkov, e-mail: mmalkov@ucsd.edu

Competing financial interests The authors declare no competing financial interests.




Figure 1: **Momentum space of accelerated protons.** Particle scattering zones on the $(p_{\parallel}, p_{\perp})$ - plane of momentum space. Particles in the stripes $p_1 < |p_{\parallel}| < p_2$ are not scattered by waves (see text). Therefore, particles from the domains $|p_{\parallel}| > p_2$ maintain their propagation direction and promptly escape from the dense gas region.




Figure 2: **Spectra of accelerated protons and electrons.** The both particle distributions are calculated for a weakly modified shock and are shown in momentum normalization ($f(p)$ is steeper by two powers than the spectra in energy normalization, used in the text). Both spectra are multiplied by p^4 , so that the test particle distribution is flat. Shock parameters: acoustic Mach number $M = 30$, shock velocity $V_s/c = 10^{-3}$, the break momentum $p_{br} \simeq 7mc$. Shock pre-compression (flow compression across the CR precursor) $R=1.8$, injection parameter $\nu \simeq 0.1$ [defined as $\nu = (4\pi/3) (mc^2/\rho V_s^2) (p_{inj}/mc)^4 f(p_{inj})$, with ρ and V_s being the ambient gas density and the shock speed, respectively]; injection momentum $p_{inj}/mc \simeq 1.4 \cdot 10^{-3}$.




Figure 3: **Gamma radiation spectra.** Photon spectra resulting from π^0 decay and calculated for two different parent proton spectra compared against the Fermi data (open circles). Solid line: a test particle acceleration regime with the spectral index $q = 2$ below the break and $q = 3$ above the break at $p_{br} = 7\text{GeV}/c$. Dashed line: a moderately nonlinear acceleration regime corresponding to the spectrum shown in Fig.2 ($q \simeq 1.75$ and $q \simeq 2.75$ below and above the break, respectively). Cut-offs are placed at 300 GeV for TP- and 100 GeV, for NL-spectrum.